

Papéis Avulsos de Zoologia

Museu de Zoologia da Universidade de São Paulo

Volume 51(20):307-323, 2011

www.mz.usp.br/publicacoes
<http://portal.revistasusp.sibi.usp.br>
www.scielo.br/paz

ISSN impresso: 0031-1049

ISSN on-line: 1807-0205

A FURTHER CONTRIBUTION TO THE KNOWLEDGE OF *PECTINIUNGUIS MINUTUS* (DEMANGE, 1968), A LITTLE KNOWN DWARF SCHENDYLID CENTIPEDE FROM WESTERN EQUATORIAL AFRICA (CHILOPODA: GEOPHILOMORPHA)

LUIS ALBERTO PEREIRA¹

ABSTRACT

The geophilomorph centipede Pectiniunguis minutus (Demange, 1968), a little known dwarf schendylid from Gabon (Western equatorial Africa), is redescribed and illustrated based on the type material and an additional non-type specimen preserved in the collections of the Muséum national d'Histoire naturelle, Paris. P. minutus can be easily differentiated from all the other species currently assigned to the genus Pectiniunguis, by the very low number of leg-bearing segments (35 in the males, 37 or 39 in the females) and very small body size (12-16 mm in length). P. minutus is also distinguished by having ventral pore-fields on the anterior region of the body only, this character being shared by a single species of the genus, i.e., P. ascendens Pereira, Minelli & Barbieri, 1994, from the Neotropics (Brazil: State of Amazonas) with which a morphological comparison is given. Comments about other dwarf centipede species belonging to several families of the order Geophilomorpha, are also added.

KEY-WORDS: *Pectiniunguis*; Taxonomy; Western Equatorial Africa; Chilopoda; Geophilomorpha; Schendylidae.

INTRODUCTION

Two hundred and nineteen species, in thirty-three genera are currently recognized in the geophilomorph family Schendylidae. The taxon occurs in the Americas including the Antilles; Europe and central-northern Asia, including the Japanese region; scattered records from Africa; Madagascar; from Indochina, through Indonesia, to Australia, New Caledonia, Hawaii and the Fiji Islands (Pereira *et al.*, 1997; Bonato *et al.*, 2009).

In sub-Saharan Africa the family currently includes twenty-three species in four genera: six species in the genus *Mesoschendyla* Attems, 1909; five

in *Schendylops* Cook, 1899; 11 in *Ctenophilus* Cook, 1896 and one in *Pectiniunguis* Bollman, 1889. The species of the first two genera are distributed in the equatorial and southernmost regions, while those of the two latter occur in the western equatorial region only. These four taxa are also widespread elsewhere: *Mesoschendyla* also occurs in Madagascar (one species) and Java (one species); *Schendylops* in Northern mainland Africa (two species), Madagascar (5 species), and the Neotropical Region (54 species); *Ctenophilus* in the Neotropics (one species); and *Pectiniunguis* in Oceania Region (one species in Fiji Islands), and the New World (21 species) distributed in southern North America (U.S.A), Central America (Mexico),

1. CONICET and Universidad Nacional de La Plata, Facultad de Ciencias Naturales y Museo, Paseo del Bosque s/n, (1900) La Plata, Argentina. E-mail: lpereira@fcnym.unlp.edu.ar

Caribbean Islands (Cuba and Cayo Sombrero (Venezuela)), South American mainland (Colombia, Guyana, continental Ecuador, Brazil and Argentina), and the Galápagos Islands. The last survey of *Mesoschendyla* can be found in Crabill 1968. Recent contributions to the study of the species of *Schendylops* from mainland Africa include Pereira & Minelli, 1995, 2001; studies on those inhabiting Madagascar comprise Hoffman & Pereira, 1997; Pereira *et al.*, 2004. The African species of *Ctenophilus* were revised by Pereira & Demange, 1991, 1997. Latest contributions to the knowledge of the Neotropical species of *Pectiniunguis* can be found in Pereira & Coscarón, 1975(1976); Pereira *et al.*, 1994, 1995, 1999, 2000, 2001; Shear & Peck, 1992; Pereira, 2010a; the single species from Fiji Islands was revised by Pereira, 1982.

The purpose of the present contribution is to redescribe the only known representative of the genus *Pectiniunguis* from Africa (Gabon), which was described by Demange (1968) under the name of *Pleuroschendyla minuta*. The original description lacks information on many important characters of specific value and only includes two inadequately detailed figures. Pereira & Demange (1991) revised the holotype and transferred the species to the genus *Pectiniunguis* on the basis of the morphological features of the pleurites of the second maxillae, but besides the proposal of the new combination [*Pectiniunguis minutus* (Demange, 1968)], no additional morphological data, nor new illustrations were given. Subsequently, no other authors have treated this taxon which has remained poorly known. The opportunity to revise all the type material and an additional non-type specimen during a visit to the Laboratoire de Zoologie (Arthropodes), Muséum national d'Histoire naturelle, Paris, allows a detailed redescription with new illustrations, giving a better understanding of this interesting dwarf schendylid centipede.

MATERIAL AND METHODS

The type and non-type material herein revised, is currently housed at the Muséum national d'Histoire naturelle, Paris (MNHN).

The specimens were examined and illustrated using a compound microscope equipped with a drawing tube attachment. Temporary mounts were prepared by direct transfer of the specimens from the preservation liquid (70 per cent ethanol) onto microscope slides, using as a clearing agent/mounting medium, undiluted 2-Phenoxyethanol (CAS No. 122-99-6). Details on preparation of microscope

slides and employed dissection procedures are described in Pereira (2000, 2010a, 2010b); Foddai *et al.* (2002). All measurements are given in mm. Terminology for the external anatomy follows Bonato *et al.* (2010, 2011). The following abbreviation was used in the text and legends of the figures: a.a., antennal article/articles.

RESULTS

Family Schendylidae

Genus *Pectiniunguis* Bollman, 1889

Diagnosis: This genus can be distinguished from other genera currently recognized in the family Schendylidae by the following particular combination of features. Second maxillae: pleurites not fused to the posterior internal border of the coxosternite, apical claw of telopodites pectinate on both dorsal and ventral edge. Sternites: with pore-fields. Ultimate leg-bearing segment: ultimate legs with seven articles, pretarsus in form of a small hairy tubercle or replaced by a small spine or altogether absent, each coxopleuron with two internal coxal organs of composite structure ("heterogeneous coxal glands" *sensu* Brölemann & Ribaut (1912)).

Type species of the genus: *Pectiniunguis americanus* Bollman, 1889, by original designation.

Species currently included in the genus: *P. albemarlensis* Chamberlin, 1913 (Ecuador: Galápagos Islands); *P. americanus* Bollman, 1889 (Ecuador: Galápagos; Mexico; U.S.A.); *P. amphibius* Chamberlin, 1923 (Mexico); *P. argentinensis* Pereira & Coscarón, 1976 (Argentina); *P. ascendens* Pereira, Minelli & Barbieri, 1994 (Brazil); *P. bolbonyx* (Brölemann & Ribaut, 1912) (Brazil); *P. bollmani* Pereira, Minelli & Foddai, 1999 (Venezuela); *P. catalinensis* Chamberlin, 1941 (U.S.A.); *P. chazaliei* (Brölemann, 1900) (Colombia); *P. ducalis* Pereira, Minelli & Barbieri, 1995 (Brazil); *P. fijiensis* (Chamberlin, 1920) (Fiji Islands); *P. gaigei* (Chamberlin, 1921) (Guyana); *P. geayi* Brölemann & Ribaut, 1911 (Brazil); *P. halirrhytus* Crabill, 1959 (Mexico; U.S.A.); *P. imperforatus* (Brölemann, 1902) (Brazil); *P. insulanus* (Brölemann & Ribaut, 1911) (Cuba); *P. krausi* Shear & Peck, 1992 (Ecuador: Galápagos Islands); *P. minutus* (Demange, 1968) (Gabon); *P. nesiotus* Chamberlin, 1923 (Mexico); *P. pauperatus* Silvestri, 1907 (West Indies); *P. pectinatus* (Attems, 1934) (Brazil); *P. plusiodontus* Attems, 1903 (Brazil); *P. roigi* Pereira, Foddai & Minelli, 2001 (Ecuador).

TABLE 1: Comparative matrix of differential morphological traits for the Schendylid genera known to occur in Sub-Saharan Africa: *Ctenophilus* Cook, 1896; *Mesoschendyla* Attems, 1909; *Pectiniunguis* Bollman, 1889 and *Schendylops* Cook, 1899.

	<i>Pectiniunguis</i>	<i>Ctenophilus</i>	<i>Schendylops</i>	<i>Mesoschendyla</i>
Pleurites of the second maxillae fused to the internal posterior border of the coxosternite	No (Figs. 14, 37-39, 45)	Yes (Fig. 44)	No	No
Coxopleura of the ultimate leg-bearing segment with 1+1 coxal organs	No	No	No	Yes
Coxopleura of the ultimate leg-bearing segment with 2+2 coxal organs	Yes	Yes	Yes	No
Coxal organs homogeneous (<i>sensu</i> Brölemann & Ribaut (1912))	No	No	Yes	Yes
Coxal organs heterogeneous (<i>sensu</i> Brölemann & Ribaut (1912))	Yes (Figs. 29, 34, 40-42; 47)	Yes	No	No

Remarks: Differential characters between *Pectiniunguis* and the other three genera of Schendylidae known to occur in Sub-Saharan Africa, *i.e.*, *Ctenophilus*, *Mesoschendyla* and *Schendylops*, are shown in Table 1.

Redescription

Pectiniunguis minutus (Demange, 1968) (Figs. 1-43)

Pleuroschendyla minuta Demange, 1968: 283-284.

Pleuroschendyla minuta: Mauriès, 1968: 771.

Pectiniunguis minutus: Pereira & Demange, 1991: 95; Pereira, Minelli & Foddai, 1999: 177.

Diagnosis: A *Pectiniunguis* species with 35, 37 or 39 leg-bearing segments; body length 12-16 mm; ventral pore-fields present on the anterior region of the body only (absent on first sternite). It can be easily differentiated from all the other species currently assigned to the genus, by the very low number of leg-bearing segments and very small body size. The following traits, are also distinctive features for this species: poison glands unusually large, extending along the entire forcipular telopodite and the anterior half of the forcipular coxosternite (Fig. 16: c); a.a. II-XIII all wider than long (Fig. 1); claw of waking legs with two accessory spines only (one anterior and one posterior, Figs. 18, 19); pleurites of second maxillae somewhat expanded on the anterior-external area (Figs. 14, 37-39); a.a. XIV with claviform sensory setae on the external edge only (Fig. 36); coxosternite of first maxillae without setae (Fig. 14); pretarsus of ultimate legs represented by a small tubercle with two diminutive apical spines (Figs. 28, 35, 43).

Type material examined: All specimens from Gabon: Bélinga, H. Coiffait leg. (Ref. *Pleuroschendyla*

minuta Demange); 25 March 1963: allotype female with 37 leg-bearing segments, body length 13 mm; 10 May 1963: holotype male with 35 leg-bearing segments, body length 12 mm; paratype male with 35 leg-bearing segments, body length 11.5 mm. (MNHN Collection Myriapodes et Onychophores: M206).

Other material examined: Gabon: Bélinga: Station 154 T, route du camp VI. Forêt tropicale, 27 February 1963, H. Coiffait leg. (Ref. *Pleuroschendyla minuta* Demange), female with 39 leg-bearing segments, body length 13 mm (MNHN Collection Myriapodes et Onychophores: M206).

Remarks: The original description mentions the four specimens cited above, and an additional non type from "Bélinga: Station 129 T, carrière, route du débarcadère, Forêt tropicale" (sex and number of leg-bearing segments not specified), but this latter was not revised in the present study.

Male paratype: Thirty-five leg-bearing segments, body length 11.5 mm, maximum body width 0.50 mm. Width of cephalic plate, *ca.* 0.41 mm. Width of forcipular coxosternite, *ca.* 0.40 mm. Ground color (of preserved specimen in alcohol) pale yellowish.

Antennae: *ca.* 2.4 times as long as the cephalic plate, distally slightly attenuate; a.a. II to XIII all wider than long, a.a. XIV longer than wide (Fig. 1). Setae on a.a. I to VI of different lengths and few in number, those of remaining articles progressively shorter and more numerous towards the tip of the appendage (Fig. 1). Apical a.a. with *ca.* 15 claviform sensilla on the external border and absent on the internal border (similar to the holotype, Fig. 36); distal end of this a.a. with *ca.* 4 specialized sensilla apparently not split apically, (similar to the holotype, Fig. 36). Dorsal and ventral

TABLE 2: Number of type *a*, *b* and *c* sensilla on a.a. II, V, IX and XIII in the male paratype of *Pectiniunguis minutus* (Demange, 1968).

	Ventral		Dorsal			Figs.
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>c</i>	
II	1	1	1	1		2
V	1	1	1	1		3, 4
IX	1	1	1	2	1	5, 6
XIII	1	1	1	1	2	7

surface of a.a. II, V, IX and XIII (Figs. 2-7) with very small specialized sensilla. On the ventral side these sensilla are restricted to an internal latero-apical area and are represented by two different types: *a* and *b*. Type *a* sensilla are very thin and not divided apically (Fig. 5: a); type *b* sensilla are similar to those on the apex of the terminal a.a. but with two very small apical branches (Fig. 5: b). Specialized sensilla on dorsal side restricted to an external latero-apical area on a.a. II, V and IX (Figs. 2, 4, 6) and to an internal and external sublateral apical areas on a.a. XIII (Fig. 7), and are represented by three different types: *a*, similar to *a* of ventral side (Fig. 7: a); *b*, similar to *b* of ventral side, with two very small apical branches on a.a. II (Fig. 2: b) but apparently not divided apically on a.a. V, IX and XIII (Figs. 4, 6-7: b); and type *c* sensilla, much bigger than type *b*, not divided apically and darker (brownish-ochre in colour) (Figs. 6-7: c). Distribution of type *a*, *b* and *c* sensilla on a.a. II, V, IX and XIII as in Table 2.

Cephalic plate: slightly longer than wide (ratio length/width *ca.* 1.10: 1), shape and chaetotaxy as in Fig. 8.

Clypeus: with 1+1 postantennal setae, 2+2 median setae and 1+1 prelabral setae arranged as in Fig. 9.

Labrum: with 18 small teeth, those of the central arc dark and round tipped, the lateral ones less sclerotized, each with a very sharp medial extension (Fig. 10).

Mandible: dentate lamella subdivided into two distinct blocks, with 5, 3 teeth; pectinate lamella with *ca.* 12 hyaline teeth (Figs. 11-12).

First maxillae: with lappets on the coxosternite and telopodites, those of coxosternite very poorly developed (Fig. 13). Coxosternite without setae; coxal projections subtriangular, well developed and provided with 1+1 setae (Fig. 14). Article II of telopodite with 2+2 setae on ventral side and 3+3 sensilla on dorsal side (Figs. 13-14).

Second maxillae: with 5+5 setae on the coxosternite arranged as in Fig. 14. Apical claw of telopodite well developed, bipectinate, ventral edge with *ca.* 12 teeth, dorsal edge with *ca.* 15 teeth (Fig. 15). Pleurites somewhat expanded on their anterior external edge (Fig. 14).

Forcipular segment: when closed, the telopodites do not extend beyond the anterior margin of the head. Forcipular tergite with an irregular transverse median row of *ca.* 12 large setae and a few additional smaller setae scattered on the remaining surface. All articles of the telopodites without teeth. Poison glands unusually large (shape and relative size as in Fig. 16: c). Calyx of poison gland subcylindrical and very short (Figs. 16, 17: b). Pilosity of coxosternite and telopodites as in Figure 16.

Walking legs: chaetotaxy similar throughout the entire body length. Distribution, number and relative size of setae as in Fig. 18. Claws with two thin and pale accessory spines ventrobasally, one anterior and one posterior (shape and relative size as in Fig. 19).

Sternites of leg-bearing segments 1 to penultimate: pore-fields present from sternite 2 to 20 inclusive (wholly absent on the remaining sternites). All fields undivided, transversally subovoidal, shape and relative size changing along the trunk as in Figs. 20-25. Number of pores on selected sternites: sternite 2 (18 pores); 4 (37); 9 (48); 14 (46); 19 (16); 20 (9).

Ultimate leg-bearing segment: intercalary pleurites absent at both sides of the ultimate pretergite. Ultimate presternite not divided along the sagittal plane; length/width ratio of the tergite, 0.73: 1; length/width ratio of the sternite, 0.56: 1. Shape and chaetotaxy of tergite and sternite as in Figs. 26-27. Coxopleura slightly protruding at their distal-internal ventral ends, setae small and numerous on the distal-internal ventral area, remaining surface with few larger setae. Two compound ('heterogeneous') coxal organs in each coxopleuron (Figs. 26, 27, 29). Coxal organs open on the membrane between coxopleuron and sternite, partially covered by the latter (Figs. 27, 29), internal cuticular structure as shown in Fig. 29 (a: individualized areas of mucous layer). Ultimate legs inflated, composed of seven articles. Ratio of length of telopodites of ultimate legs/length of sternite *ca.* 4.03: 1. Shape and chaetotaxy of ultimate legs as in Figs. 26-27. Ultimate pretarsus represented by a rudimentary sub-terminal tubercle with 2 diminutive spines (Fig. 28).

Postpedal segments: intermediate tergite with posterior margin strongly convex (Fig. 26), intermediate sternite and first genital sternite with posterior margin medially slightly convex, laterally slightly concave (Fig. 27). Gonopods biarticulate, basal article with *ca.* 10 setae, apical article with *ca.* 6 setae (Figs. 27, 30), penis with 1+1 apical setae on dorsal side (Fig. 31).

Female allotype: Thirty-seven leg-bearing segments, body length 13 mm, maximum body width 0.60 mm. All features similar to those in the male except for the shape and chaetotaxy of the ultimate leg-bearing segment and postpedal segments.

Ultimate leg-bearing segment: tergite and sternite trapézoidal, length/width ratio of tergite 0.67: 1; length/width ratio of sternite 0.63: 1. Shape and chaetotaxy of tergite and sternite as in Figs. 32-33. Coxopleura slightly protruding at their distal-internal ventral ends; setae distributed on the distal ventral-lateral areas, the remaining surface without setae (Figs. 32-33). Articles of ultimate legs not inflated, much thinner than those of the male (Figs. 32-33). Ultimate legs proportionally much shorter than those of the male, with ratio of length of telopodites/length of sternite *ca.* 2.11: 1. Shape and chaetotaxy of ultimate legs as in Figs. 32-33.

Postpedal segments: intermediate tergite with posterior margin strongly convex (Fig. 32), intermediate sternite with posterior margin straight to slightly concave; first genital sternite with posterior margin medially convex, laterally slightly concave (Fig. 33). Gonopods uniarticulate (Fig. 33).

Variation: males with 35, and females with 37 or 39 leg-bearing segments.

The posterior limit of the ventral pore-field series is at the sternite 20 in the male holotype and male paratype (both with 35 leg-bearing segments); sternite 22 in the female allotype (with 37 leg-bearing segments); and sternite 28 in the non-type female (with 39 leg-bearing segments).

Anterior coxal organs with two outer lobes, posterior coxal organs with two-three outer lobes (Fig. 29: b); each lobe internally, with one-three individualized areas of mucous layer (Figs. 29: a, 34, 40-42). (For details on fine structure and function of coxal organs, *see* Rosenberg & Seifert (1977); Lewis (1981); Rosenberg (1982, 1983)).

The body is (according to the original description) “12-16 mm” long. (View comments on this trait, below).

Other features with non-significant variation.

Remarks: The adult condition of the four specimens herein revised is indicated by the tubula seminifera full of mature spermatozoa in the males, and the two spermathecae full of spermatozoa in the females.

The original description by Demange is very short, only includes two figures (labrum, and ultimate leg-bearing segment and postpedal segments in ventral view), and completely lacks information on chaetotaxy of the antennae; kind and number of specialized sensilla on a.a. II, V, IX and XIII; pilosity of clypeus; shape and chaetotaxy of the forcipular segment; relative size of the ventral pore-fields; pilosity of walking legs; details of internal structure of the coxal organs; form of pretarsus of the ultimate legs; shape of the female postpedal segments; etc.


The author states “champs poreux présents du 2e segment au 21-22e segment” (but the posterior limit of the ventral pore-field series varies as detailed above). Of the body length he says “12-16 mm”, but none of the four specimens here revised exceeds 13 mm in length, “16 mm” could refer to the specimen from “Bélinga. Station 29 T, carrière, route du débarcadère. Forêt tropicale” (not examined in the present contribution, as stated above).

Type locality: Gabon: Bélinga region.


Known range: Only known from the type locality.

DISCUSSION


The original description of *Pectiniunguis minutus* mentions as type locality the Bélinga region in Gabon (which according to White (1983) is part of the Guineo-Congolian rainforest). No data are given on the collecting sites of the three type specimens, however it is stated for the two non type specimens that they come from sites in a tropical forest environment (but the altitude a.s.l. is not given). In western equatorial Africa, the tropical lowland rainforest of Gabon is well known for its high biodiversity (Sosef 1994); according to Ngomanda *et al.* (2009) this forest is today well conserved because of the absence of intensive agricultural activities, coupled with low population densities. (Nevertheless, the biodiversity in the Bélinga mountains



FIGURES 1-5: *Pectiniunguis minutus* (Demange, 1968), (male paratype; Gabon: Bélinga): (1) Left antenna, ventral; (2) Left a.a. II, dorsal (a, b: a, b type sensilla); (3) Left a.a. V, ventral (a, b: a, b type sensilla); (4) Left a.a. V, dorsal (a, b: a, b type sensilla); (5) Left a.a. IX, ventral (a, b: a, b type sensilla). Scale bars: 0.05 mm (2-5); 0.2 mm (1).



FIGURES 6-12: *Pectiniunguis minutus* (Demange, 1968), (male paratype; Gabon: Bélinga): (6) Left a.a. IX, dorsal (a, b, c: a, b, c type sensilla); (7) Left a.a. XIII, dorsal (a, b, c: a, b, c type sensilla); (8) Cephalic plate and bases of antennae; (9) Clypeus and bases of antennae; (10) Labrum; (11) Right mandible; (12) Left mandible. Scale bars: 0.05 mm (6-7, 10-12); 0.2 mm (8-9).




FIGURES 13-19: *Pectiniunguis minutus* (Demange, 1968), (male paratype; Gabon: Bélinga): (13) Left side of first maxillae, dorsal; (14) First and second maxillae, ventral; (15) Claw of right telopodite of second maxillae, dorsal; (16) Forcipular segment, ventral (a: duct, b: calyx, c: poison gland); (17) Detail of the duct (a) and calyx (b) of poison gland in left forcipular telopodite, ventral; (18) Left waking leg (pair 8), ventral; (19) Claw of left waking leg (pair 8), ventral view. Scale bars: 0.03 mm (15, 19); 0.1 mm (13, 17-18); 0.2 mm (14, 16).

is threatened by global warming and iron mining (Leal, 2008)).

P. minutus is characterized by having thin and pale accessory spines on the claws of the walking legs (Fig. 19), this trait is shared by most of the species of *Pectiniunguis* inhabiting South America and the one from the Fiji Islands. (The opposite condition, *i.e.*, strong and dark accessory spines, occur in some members distributed in southern North America, Central America, West Indies and Northwestern South America, *e.g.*, *P. bollmani* Pereira, Minelli & Foddai, 1999 from Venezuela (Fig. 48)). Among the species currently recognized in the genus, only *P. ascendens* Pereira, Minelli & Barbieri, 1994, from the


Neotropical Region (Brazil: State of Amazonas) shares with *P. minutus* the peculiarity of bearing ventral pore-fields on the anterior region of the body only. These two species also share a similar shape of pretarsus of the ultimate legs (represented by a rudimentary tubercle with 2 diminutive spines); a similar internal chitinous structure of the coxal organs (showing one to three individualized areas of mucous layer for each outer lobe); presence of specialized brownish-ochre sensilla, on dorsal side of some a.a. A morphological comparison of these apparently similar taxa, can be made by means of the following selected traits (those for *P. ascendens* are given in parentheses): males with 35, females with 37 or 39 leg-bearing segments




FIGURES 20-25: *Pectiniunguis minutus* (Demange, 1968), (male paratype; Gabon: Bélinga): (20) Sternite of leg-bearing segment 2; (21) Sternite of leg-bearing segment 4; (22) Sternite of leg-bearing segment 9; (23) Sternite of leg-bearing segment 14; (24) Sternite of leg-bearing segment 19; (25) Sternite of leg-bearing segment 20. Scale bar: 0.1 mm.

(males with 43 or 45, females with 43, 45 or 47); maximum body length 16 mm (33 mm); a.a. II–XIII all wider than long, Fig. 1 (all longer than wide); specialized brownish-ochre sensilla present on dorsal side of a.a. IX and XIII (on dorsal side of a.a. V, IX and XIII); clypeus with *ca.* 4 setae on the middle,


Fig. 9 (with *ca.* 16 setae); pleurites of second maxillae somewhat expanded on the anterior external area, Figs. 14, 37–39 (pleurites not expanded on the anterior external area, Fig. 45); lappets of coxites of first maxillae poorly developed, Fig. 13 (well-developed); dentate lamellae of mandibles divided in two blocks,




FIGURES 26–28: *Pectiniunguis minutus* (Demange, 1968), (male paratype; Gabon: Bélinga): **(26)** Ultimate leg-bearing segment and postpedal segments, dorsal; **(27)** Ultimate leg-bearing segment and postpedal segments, ventral; **(28)** Detail of distal end of apical article of right ultimate leg, ventral (a: pretarsus). Scale bars: 0.02 mm (28); 0.4 mm (26–27).



FIGURES 29-32: (29-31) *Pectiniunguis minutus* (Demange, 1968), (male paratype; Gabon: Bélinga): (29) Left coxal organs, ventral (a: individualized areas of mucous layer, b: outer lobe); (30) Left gonopod, ventral; (31) Penis, dorsal. (32) *Pectiniunguis minutus* (Demange, 1968), (female allotype; Gabon: Bélinga): Ultimate leg-bearing segment and postpedal segments, dorsal. Scale bars: 0.04 mm (30-31); 0.1 mm (29); 0.2 mm (32).




FIGURES 33-38: (33-35) *Pectiniunguis minutus* (Demange, 1968), (female allotype; Gabon: Bélinga): (33) Ultimate leg-bearing segment and postpedal segments, ventral; (34) Left coxal organs, ventral (a: individualized areas of mucous layer, b: outer lobe); (35) Detail of distal end of apical article of left ultimate leg, ventral (a: pretarsus). (36-38) *Pectiniunguis minutus* (Demange, 1968), (male holotype; Gabon: Bélinga): (36) Distal half of right a.a. XIV, ventral view (a: claviform sensilla, b: apical specialized sensilla); (37) Detail of pleurite, coxopodite and base of telopodite of second maxillae, right side, ventral (a: pleurite); (38) Pleurite, coxopodite and base of telopodite of second maxillae, left side, dorsal (a: pleurite). Scale bars: 0.03 mm (35); 0.05 mm (36); 0.1 mm (34, 37-38); 0.2 mm (33).



FIGURES 39-43: (39) *Pectiniunguis minutus* (Demange, 1968), (male holotype; Gabon: Bélinga): Detail of pleurite, coxopodite and basis of telopodite of second maxillae, left side, ventral (a: pleurite). (40-43) *Pectiniunguis minutus* (Demange, 1968), (female non type; Gabon: Bélinga: Station 154 T, route du camp VI. Forêt tropicale): (40) Right coxal organs, ventral (a: individualized areas of mucous layer); (41) Left coxal organs, ventral (a: individualized areas of mucous layer, b: outer lobe); (42) Coxal organs, dorsal (a: individualized areas of mucous layer); (43) Detail of distal end of apical article of right ultimate leg, ventral (a: pretarsus). Scale bars: 0.02 mm (43); 0.1 mm (39-42).

Figs. 11-12 (divided in three blocks); claw of walking legs with two accessory spines, one anterior and one posterior, Fig. 19 (with three accessory spines, one anterior and two posterior, Fig. 46). (The precedent

comparison, merely express similarities and differences between presumptively morphologically similar species. Phylogenetic relationships can only be elucidate through appropriate analyses).



FIGURES 44-48: (44) *Ctenophilus amieti* (Demange, 1963), (reference *Pleuroschendyla magna*, female holotype; Ivory Coast: Mont Nimba): Detail of pleurite, coxopodite and basis of telopodite of second maxillae, right side, ventral (a: pleurite). (After Pereira & Demange 1997). (45-47) *Pectiniunguis ascendens* Pereira, Minelli & Barbieri, 1994, (female holotype; Brazil: Amazonas: Rio Trumã Mirim): (45) Detail of pleurite and proximal external portion of coxopodite of second maxillae, left side, ventral (a: pleurite); (46) Claw of left walking leg (pair 12), antero-ventral view; (47) Detail of left coxal organs, ventral (a: individualized area of mucous layer, b: outer lobe). (After Pereira *et al.*, 1994). (48) *Pectiniunguis bollmani* Pereira, Minelli & Foddai, 1999 (male holotype; Venezuela: State of Falcón: Parque Nacional Morrocoy: Cayo Sombrero): Claw of right walking leg (pair 60), ventral. (After Pereira *et al.*, 1999). Scale bars: 0.03 mm (46); 0.05 mm (48); 0.1 mm (45, 47); no scale available (44).

As mentioned above, *Pectiniunguis minutus* was originally placed in the genus *Pleuroschendyla* Brölemann & Ribaut, 1911, for which the current valid name is *Ctenophilus* Cook, 1896 (senior synonym of the first). Twelve species are currently included in *Ctenophilus*; *Pectiniunguis minutus* can be easily differentiated from all of them by the shape of the pleurites of the second maxillae which are not fused to the posterior internal border of the coxosternite (Figs. 14, 37-39). (In all species at present assigned to *Ctenophilus*, those pleurites are fused to the postero-internal area of the second maxillary coxosternite, as shown in Fig. 44). At least five African species of *Ctenophilus* share with *P. minutus* the particularity of having pore-fields on the anterior region of the body only: *C. angolae* (Chamberlin, 1951) (from Zaire: Gombe Louzi, Thysville; Angola: Dundo); *C. corticeus* (Demange, 1968) (from Gabon: Bélinga); *C. edentulus* (Porat, 1894) (from Camerun: Bonge, Olounou; Gabon: Makokou, Bélinga); *C. nitidus* (Brölemann, 1926) (from Bénin: Athiéme) and *C. oligopodus* (Demange, 1963) (from Ivory Coast: Mont Nimba). (But in its original description, *P. minutus* is compared with the latter only).

As *Pectiniunguis minutus*, diverse other cases of reduced body size exist within geophilomorphs (see Minelli *et al.* 2000, Foddai *et al.* 2003, Pereira 2009). This is known to occur in some genera of the Geophilidae (within *Geophilus* Leach, 1814, where *G. persephones* Foddai & Minelli, 1999, with 29 pairs of legs is 16 mm long, *Hyphydrophilus* Pereira, Minelli & Barbieri, 1994, *Ribautia* Brölemann, 1909, and *Dinogeophilus* Silvestri, 1919, where *D. oligopodus* Pereira, 1984, with 29 pairs of legs, is just 4.5 mm long); Linotaenidae (within *Strigamia* Gray, 1893, where *S. hoffmani* Pereira, 2009, with 33, 35 or 37 leg-bearing segments is 12-16 mm long); the Schendylidae (within *Schendyla* Bergsøe and Meinert, 1866 and *Schendylops* Cook, 1899, where *Schendylops oligopus* (Pereira, Minelli & Barbieri, 1995), with 27, 29, and 31 pairs of legs is 8-9 mm long); the Ballophilidae (within *Taeniolum* Pocock, 1893, where *T. arborum* Pereira, Minelli & Barbieri, 1994, with 43 or 45 leg-bearing segments is 10-13 mm long); and the Mecistocephalidae (within *Nannarrup* Foddai, Bonato, Pereira & Minelli, 2003, where *N. hoffmani* Foddai, Bonato, Pereira & Minelli, 2003, with 41 pairs of legs is 10 mm long). Reduction in adult body size hence evolved independently in several derived lineages, often coupled with a secondary reduction to the smallest number of pairs of legs actually found

in the respective family; nevertheless, no reduction in the number of body segments is involved in some species such as *N. hoffmani*, which was analysed as a possible instance of miniaturization (Foddai *et al.*, 2003).

It remains to be seen whether the numerous species of Schendylidae currently referred to the genus *Pectiniunguis* really form a monophyletic unit. Their geographical distribution with a majority of species in the New World, accompanied by a single one on mainland Africa, and one in the Oceania region, invites closer investigation; but a phylogenetical study is not an easy task as the whole generic classification of schendylid geophilomorphs is still based on characters of uncertain phylogenetic value.

Only three species of Schendylidae (in two genera) are known to occur in Gabon: *Ctenophilus corticeus* (Demange, 1968), *Ctenophilus edentulus* (Porat, 1894) and *Pectiniunguis minutus* (Demange, 1968). They may only represent a small portion of the biodiversity of the family in the rich tropical rainforest biome of that country. It is possible that intensive field work on the whole extension of the region, may result in the discovery of new taxa of geophilomorph centipedes.

RESUMEN

El quilópodo geofilomorfo Pectiniunguis minutus (Demange, 1968), una especie enana de Schendylidae pobremente conocida de Gabon (Africa ecuatorial occidental), se redescubre e ilustra en base al material tipo y a un espécimen adicional no tipo conservados en las colecciones del Muséum national d'Histoire naturelle, París. *P. minutus* se puede diferenciar fácilmente de todas las otras especies actualmente asignadas al género *Pectiniunguis*, por el número muy bajo de segmentos pedales (35 en los machos, 37 o 39 en las hembras) y la pequeña longitud del cuerpo (12-16 mm). *P. minutus* también se distingue por tener campos de poros ventrales solamente en la región anterior del cuerpo, carácter que es compartido por una sola especie del género, i.e., *P. ascendens* Pereira, Minelli & Barbieri, 1994, distribuida en la Región Neotropical (Brasil: Estado de Amazonas), con la cual una comparación morfológica es brindada. Comentarios sobre otras especies enanas de ciempiés pertenecientes a diversas familias del orden Geophilomorpha, son también incluidos.

PALABRAS-CLAVE: *Pectiniunguis*; Taxonomía; Africa Equatorial Occidental; Chilopoda; Geophilomorpha; Schendylidae.

ACKNOWLEDGEMENTS

I am indebted to Jean-Paul Mauriès, Monique Nguyen Duy-Jacquemin, Jean-Marie Demange and Jean-Jacques Geoffroy of the Muséum national d'Histoire naturelle, Paris for the access to relevant specimens, library facilities, valuable guidance and hospitality in the Laboratoire de Zoologie (Arthropodes) during part of this study. I am also grateful to the Museum authorities for the financial support that enabled my stay at the institution. Miguel E. Leal and James C. Solomon (Missouri Botanical Garden, St. Louis), provided information on environmental features and biodiversity of the Bélinga mountains, Gabon. Two anonymous referees contributed with careful reviews which helped to improve the final version of the manuscript. Carlos José Einicker Lamas (Museu de Zoologia da Universidade de São Paulo), as well as Adriana Nascimento Flamino and Airton de Almeida Cruz (of the same institution), very kindly assisted me during all steps of the review and publication process. Hernán Lucas Pereira (La Plata) prepared the electronic versions of the figures.

REFERENCES

- BONATO, L.; BEVILACQUA, S. & MINELLI, A. 2009. An outline of the geographical distribution of world Chilopoda. *Contributions to Natural History*, Bern, 12:183-209.
- BONATO, L.; DRAGO, L. & MINELLI, A. 2011. Pincer-like claws in centipedes (Chilopoda): multiple evolutionary origin of similar form and serial pattern. *Zoomorphology*, 130:17-29.
- BONATO, L.; EDGEcombe, G.D.; LEWIS, J.G.E.; MINELLI, A.; PEREIRA, L.A.; SHELLEY, R.M. & ZAPPAROLI, M. 2010. A common terminology for the external anatomy of centipedes (Chilopoda). *Zookeys*, 69:17-51.
- BRÖLEMAN, H.W. & RIBAUT, H. 1912. Essai d'une monographie des Schendylina (Myriapodes, Géophilomorphes). *Nouvelles Archives du Muséum national d'Histoire naturelle, Paris, Sér. 5*, 4:53-183.
- CRABILL JR., R.E. 1968. Two new species of *Mesoschendyla* from the old world tropics, with a key to their congeners. *Revue de Zoologie et de Botanique africaines*, 77(3-4):283-288.
- DEMANGE, J.-M. 1968. Myriapodes Chilopodes du Gabon. *Biologia Gabonica*, 4(3):281-294.
- FODDAI, D.; BONATO, L.; PEREIRA, L.A. & MINELLI, A. 2003. Phylogeny and systematics of the Arrupinae (Chilopoda Geophilomorpha Mecistocephalidae) with the description of a new dwarfed species. *Journal of Natural History*, 37:1247-1267.
- FODDAI, D.; MINELLI, A. & PEREIRA, L.A. 2002. Chilopoda Geophilomorpha. In: Adis, J. (Ed.), *Amazonian Arachnida & Myriapoda*. Pensoft Publ., Sofia, Moscow, p. 459-474.
- HOFFMAN, R.L. & PEREIRA, L.A. 1997. The identity and taxonomic status of the generic names *Schendyllops* Cook, 1899, and *Schendylurus* Silvestri, 1907, and the proposal of *Orygmadyla*, a new related genus from Peru (Chilopoda: Geophilomorpha: Schendylidae). *Myriapodologica*, 5(2):9-32.
- LEAL, M.E. 2008. The biodiversity of the Belinga Mountains. 10 pp. Available at: <http://carpe.umd.edu/resources/Documents/BelingasMts_MELeal_Mar2008.pdf>.
- LEWIS, J.G.E. 1981. *The Biology of centipedes*. Cambridge University Press, Cambridge, 476 pp.
- MAURIÈS, J.-P. 1968. Myriapodes de l'Île aux singes (Makokou, Gabon) récoltés par J.A. Barra. *Bulletin du Muséum national d'Histoire naturelle, Paris, Série 2*, 40(4):768-771.
- MINELLI, A.; FODDAI, D.; PEREIRA, L.A. & LEWIS, J.G.E. (2000). The evolution of segmentation of centipede trunk and appendages. *Journal of Zoological Systematics and Evolutionary Research*, 38:103-117.
- NGOMANDA, A.; CHEPSTOW-LUSTY, A.; MAKAYA, M.; FAVIER, C.; SCHEVIN, P.; MALEY, J.; FONTUGNE, M.; OSLISLY, R. & JOLLY, D. 2009. Western equatorial African forest-savanna mosaics: a legacy of late Holocene climatic change? *Climate of the Past*, 5:647-659.
- PEREIRA, L.A. 1982. Nuevos aportes al conocimiento de *Pectiniunguis fijiensis* (Chamberlin, 1920). (Chilopoda: Geophilomorpha: Schendylidae). *Revista de la Sociedad Entomologica Argentina*, 41(1-4):147-153.
- PEREIRA, L.A. 2000. The preparation of centipedes for microscopical examination with particular reference to the Geophilomorpha. *Bulletin of the British Myriapod Group*, 16:22-25.
- PEREIRA, L.A. 2009. A New Dwarf Species of the Genus *Strigamia* Gray, 1843 from the Southern Appalachian Mountains of Western Virginia (Chilopoda: Geophilomorpha: Linotaeniidae). In: Roble, S.M. & Mitchell J.C. (Eds.), *A Lifetime of Contributions to Myriapodology and the Natural History of Virginia: A Festschrift in Honor of Richard L. Hoffman's 80th Birthday*. Virginia Museum of Natural History, Martinsville, VA, p. 209-222. (Special Publication No. 16).
- PEREIRA, L.A. 2010a. Redescription of *Pectiniunguis gaigei* (Chamberlin, 1921) from Guyana, with new distributional data and complementary descriptive notes on similar Neotropical species (Chilopoda: Geophilomorpha: Schendylidae). *Papéis Avulsos de Zoologia*, 50(20):297-321.
- PEREIRA, L.A. 2010b. First record of a ballophilid centipede from French Guiana with a description of *Ityphilus bettschi* sp. nov. (Myriapoda: Chilopoda: Geophilomorpha). *Papéis Avulsos de Zoologia*, 50(42):643-665.
- PEREIRA, L.A. & COSCARÓN, S. 1975(1976). Estudios sobre Geofilomorfos neotropicales I. Sobre dos especies nuevas del género *Pectiniunguis* Bollman (Schendylidae-Chilopoda). *Revista de la Sociedad Entomologica Argentina*, 35:59-75.
- PEREIRA, L.A. & DEMANGE, J.-M. 1991. Sur les espèces du genre *Ctenophilus* Cook, 1896 (Chilopoda, Geophilomorpha, Schendylidae). *Bulletin du Muséum national d'Histoire naturelle, Paris, Série 4*, 13A(1-2):81-96.
- PEREIRA, L.A. & DEMANGE, J.-M. 1997. Nouvelle contribution à la connaissance du genre *Ctenophilus* Cook, 1896, à repartition géographique disjointe (Myriapoda, Chilopoda, Geophilomorpha, Schendylidae). *Zoosystema*, 19(2-3):293-326.
- PEREIRA, L.A. & MINELLI, A. 1995. The African species of the genus *Schendylurus* Silvestri, 1907 (Chilopoda: Geophilomorpha: Schendylidae). *Memorie della Società Entomologica Italiana*, 73:29-58.
- PEREIRA, L.A. & MINELLI, A. 2001. A redescription of the South African centipede *Schendyllops caledonicus* (Attems, 1928) (Chilopoda Geophilomorpha Schendylidae). *Bolletino della Società Entomologica Italiana*, Genova, 133(1):13-25.
- PEREIRA, L.A.; FODDAI, D. & MINELLI, A. 1997. Zoogeographical aspects of Neotropical Geophilomorpha. *Entomologica Scandinavica*, Supplement, 51:77-86.
- PEREIRA, L.A.; FODDAI, D. & MINELLI, A. 2000. New taxa of Neotropical Geophilomorpha (Chilopoda). *Amazoniana*, 16(1/2):1-57.
- PEREIRA, L.A.; FODDAI, D. & MINELLI, A. 2001. *Pectiniunguis roigi* n. sp., from the Amazonian rainforest of Ecuador (Chilopoda:

- Geophilomorpha: Schendylidae). *Studies on Neotropical Fauna & Environment*, 36(2):143-150.
- PEREIRA, L.A.; MINELLI, A. & BARBIERI, F. 1994. New and little known geophilomorph centipedes from Amazonian inundation forests near Manaus, Brazil (Chilopoda: Geophilomorpha). *Amazoniana*, 13(1/2):163-204.
- PEREIRA, L.A.; MINELLI, A. & BARBIERI, F. 1995. Description of nine new centipede species from Amazonia and related matters on Neotropical geophilomorphs (Chilopoda: Geophilomorpha). *Amazoniana*, 13(3/4):325-416.
- PEREIRA, L.A.; MINELLI, A. & FODDAI, D. 1999. *Pectiniunguis bollmani* n. sp., from the coralline island Cayo Sombrero (Venezuela) with notes on *P. halirrhytus* Crabill, 1959 (Chilopoda: Geophilomorpha: Schendylidae). *Studies on Neotropical Fauna & Environment*, 34:176-185.
- PEREIRA, L.A.; MINELLI, A. & ULIANA, M. 2004. The species of *Schendyllops* Cook, 1899 (Chilopoda, Geophilomorpha, Schendylidae) from Madagascar. *Zoosystema*, 26(4):727-752.
- ROSENBERG, J. 1982. Coxal organs in Geophilomorpha (Chilopoda). Organization and fine structure of the transporting epithelium. *Zoomorphology*, 100:107-120.
- ROSENBERG, J. 1983. Coxal organs of *Lithobius forficatus* (Myriapoda, Chilopoda). Fine-structural investigation with special reference to the transport epithelium. *Cell and Tissue Research*, 230:421-430.
- ROSENBERG, J. & SEIFERT, G. 1977. The coxal glands of Geophilomorpha (Chilopoda): organs of osmoregulation. *Cell and Tissue Research*, 182:247-251.
- SHEAR, W.A. & PECK, S.B. 1992. Centipeds (Chilopoda) and Symphyla of the Galapagos Islands, Ecuador. *Canadian Journal of Zoology*, 70:2260-2274.
- SOSEF, M.S.M. 1994. Refuge Begonias: Taxonomy, phylogeny and historical biogeography of Begonia sect. Loasibegonia and sect. Scutobegonia in relation to glacial rain forest refuges in Africa. *Wageningen Agricultural University Papers*, 94:1-306. Studies in Begoniaceae, 5.
- WHITE, F. 1983. *The vegetation of Africa*, a descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa. Unesco, Paris, p. 356. (Natural Resources Research, 20)

Recebido em: 12.04.2011

Aceito em: 28.07.2011

Impresso em: 30.09.2011

EDITORIAL COMMITTEE

Publisher: Museu de Zoologia da Universidade de São Paulo. Avenida Nazaré, 481, Ipiranga, CEP 04263-000, São Paulo, SP, Brasil.

Editor-in-Chief: Carlos José Einicker Lamas, Serviço de Invertebrados, Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42.494, CEP 04218-970, São Paulo, SP, Brasil. E-mail: editormz@usp.br.

Associate Editors: Mário César Cardoso de Pinna (Museu de Zoologia, Universidade de São Paulo, Brasil); Luís Fábio Silveira (Museu de Zoologia, Universidade de São Paulo, Brasil); Marcos Domingos Siqueira Tavares (Museu de Zoologia, Universidade de São Paulo, Brasil); Sérgio Antonio Vanin (Instituto de Biotecnologia, Universidade de São Paulo, Brasil); Hussam El Dine Zaher (Museu de Zoologia, Universidade de São Paulo, Brasil).

Editorial Board: Aziz Nacib Ab'Saber (Universidade de São Paulo, Brasil); Rüdiger Bieler (Field Museum of Natural History, U.S.A.); Walter Antonio Pereira Boeger (Universidade Federal do

Paraná, Brasil); Carlos Roberto Ferreira Brandão (Universidade de São Paulo, Brasil); James M. Carpenter (American Museum of Natural History, U.S.A.); Ricardo Macedo Corrêa e Castro (Universidade de São Paulo, Brasil); Matéo de Vivo (Universidade de São Paulo, Brasil); Marcos André Raposo Ferreira (Museu Nacional, Rio de Janeiro, Brasil); Darrel R. Frost (American Museum of Natural History, U.S.A.); William R. Heyer (National Museum of Natural History, U.S.A.); Ralph W. Holzenthal (University of Minnesota, U.S.A.); Adriano Brilhante Kury (Museu Nacional, Rio de Janeiro, Brasil); Gerardo Lamas (Museo de Historia Natural "Javier Prado", Lima, Peru); John G. Maisey (American Museum of Natural History, U.S.A.); Naércio Aquino Menezes (Universidade de São Paulo, Brasil); Christian de Muizon (Muséum National d'Histoire Naturelle, Paris, France); Nelson Papavero (Universidade de São Paulo, Brasil); James L. Patton (University of California, Berkeley, U.S.A.); Richard O. Prum (University of Kansas, U.S.A.); Olivier Rieppel (Field Museum of Natural History, U.S.A.); Miguel Trefaut Urbano Rodrigues (Universidade de São Paulo, Brasil); Randall T. Schuh (American Museum of Natural History, U.S.A.); Ubirajara Ribeiro Martins de Souza (Universidade de São Paulo, Brasil); Paulo Emílio Vanzolini (Universidade de São Paulo, Brasil); Richard P. Vari (National Museum of Natural History, U.S.A.).

INSTRUCTIONS TO AUTHORS - (April 2007)

General Information: *Papéis Avulsos de Zoologia* (PAZ) and *Arquivos de Zoologia* (AZ) cover primarily the fields of Zoology, publishing original contributions in systematics, paleontology, evolutionary biology, ontogeny, faunistic studies, and biogeography. *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* also encourage submission of theoretical and empirical studies that explore principles and methods of systematics.

All contributions must follow the International Code of Zoological Nomenclature. Relevant specimens should be properly curated and deposited in a recognized public or private, non-profit institution. Tissue samples should be referred to their voucher specimens and all nucleotide sequence data (aligned as well as unaligned) should be submitted to GenBank (www.ncbi.nih.gov/Genbank) or EMBL (www.ebi.ac.uk).

Peer Review: All submissions to *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* are subject to review by at least two referees and the Editor-in-Chief. All authors will be notified of submission date. Authors may suggest potential reviewers. Communications regarding acceptance or rejection of manuscripts are made through electronic correspondence with the first or corresponding author only. Once a manuscript is accepted providing changes suggested by the referees, the author is requested to return a revised version incorporating those changes (or a detailed explanation of why reviewer's suggestions were not followed) within fifteen days upon receiving the communication by the editor.

Proofs: Page-proofs with the revised version will be sent to e-mail the first or corresponding author. Page-proofs *must be returned to the editor, preferentially within 48 hours*. Failure to return the proof promptly may be interpreted as approval with no changes and/or may delay publication. Only necessary corrections in proof will be permitted. Once page proof is sent to the author, further alterations and/or significant additions of text are permitted only at the author's expense or in the form of a brief appendix (note added in proof).

Submission of Manuscripts: Manuscripts should be sent to the **SciELO Submission** (<http://submission.scielo.br/index.php/paz/login>), along with a submission letter explaining the importance and originality of the study. Address and e-mail of the corresponding author must be always updated since it will be used to send the 50 reprints in titled by the authors. Figures, tables and graphics **should not** be inserted in the text. Figures and graphics should be sent in separate files with the following formats: ".JPG" and ".TIF" for figures, and ".XLS" and ".CDR" for graphics, with 300 DPI of minimum resolution. Tables should be placed at the end of the manuscript.

Manuscripts are considered on the understanding that they have not been published or will not appear elsewhere in substantially the same or abbreviated form. The criteria for acceptance of articles are: quality and relevance of research, clarity of text, and compliance with the guidelines for manuscript preparation.

Manuscripts should be written preferentially in English, but texts in Portuguese or Spanish will also be considered. Studies with a broad coverage are encouraged to be submitted in English. All manuscripts should include an abstract and key-words in English and a second abstract and key-words in Portuguese or Spanish.

Authors are requested to pay attention to the instructions concerning the preparation of the manuscripts. Close adherence to the guidelines will expedite processing of the manuscript.

Manuscript Form: Manuscripts should not exceed 150 pages of double-spaced, justified text, with size 12 and source Times New Roman (except for symbols). Page format should be A4 (21 by 29.7 cm), with 3 cm of margins. The pages of the manuscript should be numbered consecutively.

The text should be arranged in the following order: **Title Page, Abstracts with Key-Words, Body of Text, Literature Cited, Tables, Appendices, and Figure Captions**. Each of these sections should begin on a new page.

(1) **Title Page:** This should include the **Title, Short Title, Author(s) Name(s) and Institutions**. The title should be concise and, where appropriate, should include mention of families and/or higher taxa. Names of new taxa should not be included in titles.

(2) **Abstract:** All papers should have an abstract in **English** and another in **Portuguese or Spanish**. The abstract is of great importance as it may be reproduced elsewhere. It should be in a form intelligible if published alone and should summarize the main facts, ideas, and conclusions of the article. Telegraphic abstracts are strongly discouraged. Include all new taxonomic names for referencing purposes. Abbreviations should be avoided. It should not include references. Abstracts and key-words should not exceed 350 and 5 words, respectively.

(3) **Body of Text:** The main body of the text should include the following sections: **Introduction, Material and Methods, Results, Discussion, Conclusion, Acknowledgments, and References at end**. Primary headings in the text should be in capital letters, in bold and centered. Secondary headings should be in capital and lower case letters, in bold and centered. Tertiary headings should be in capital and lower case letters, in bold and indented at left. In all the cases the text should begin in the following line.

(4) **Literature Cited:** Citations in the text should be given as: Silva (1998) *or* Silva (1998:14-20) *or* Silva (1998: figs. 1, 2) *or* Silva (1998a, b) *or* Silva & Oliveira (1998) *or* (Silva, 1998) *or* (Rangel, 1890; Silva & Oliveira, 1998a, b; Adams, 2000) *or* (Silva, *pers. com.*) *or* (Silva *et al.*, 1998), the latter when the paper has three or more authors. The reference need not be cited when authors and date are given only as authority for a taxonomic name.

(5) **References:** The literature cited should be arranged strictly alphabetically and given in the following format:

- **Journal Article** - Author(s). Year. Article title. *Journal name*, volume: initial page-final page. Names of journals must be spelled out in full.
- **Books** - Author(s). Year. *Book title*. Publisher, Place.
- **Chapters of Books** - Author(s). Year. Chapter title. *In*: Author(s) ou Editor(s), *Book title*. Publisher, Place, volume, initial page-final page.
- **Dissertations and Theses** - Author(s). Year. *Dissertation title*. (Ph.D. Dissertation). University, Place.
- **Electronic Publications** - Author(s). Year. *Title*. Available at: <electronic address>. Access in: date.

Tables: All tables must be numbered in the same sequence in which they appear in text. Authors are encouraged to indicate where the tables should be placed in the text. They should be comprehensible without reference to the text. Tables should be formatted with vertical (portrait), not horizontal (landscape), rules. In the text, tables should be referred as Table 1, Tables 2 and 4, Tables 2-6. Use "TABLE" in the table heading.

Illustrations: Figures should be numbered consecutively, in the same sequence that they appear in the text. Each illustration of a composite figure should be identified by capital letters and referred in the text as: Fig. 1A, Fig. 1B, for example. When possible, letters should be placed in the left lower corner of each illustration of a composite figure. Hand-written lettering on illustrations is unacceptable. Figures should be mounted in order to minimize blank areas between each illustration. Black and white or color photographs should be digitized in high resolution (300 DPI at least). Use "Fig(s)." for referring to figures in the text, but "FIGURE(S)" in the figure captions and "fig(s)." when referring to figures in another paper.

Responsability: Scientific content and opinions expressed in this publication are sole responsibility of the respective authors.

Copyrights: The journals *Papéis Avulsos de Zoologia* and *Arquivos de Zoologia* are licensed under a Creative Commons Licence (<http://creativecommons.org>).

For other details of manuscript preparation of format, consult the CBE Style Manual, available from the Council of Science Editors (www.councilscienceeditors.org/publications/style).

Papéis Avulsos de Zoologia and *Arquivos de Zoologia* are publications of the Museu de Zoologia da Universidade de São Paulo (www.mz.usp.br). Always consult the Instructions to Authors printed in the last issue or in the electronic home pages: www.scielo.br/paz or www.mz.usp.br/publicacoes.